

- reflux. Endovenous Reflux Management Study Group. *J Vasc Surg* 2001;32:330-42.
12. Dwerryhouse S, Davies B, Harradine K, Earnshaw JJ. Stripping the long saphenous vein reduces the rate of reoperation for recurrent varicose veins: five-year results of a randomized trial. *J Vasc Surg* 1999;29:589-92.
  13. Fischer R, Linde N, Duff C, Jeanneret C, Chandler JG, Seeber P. Late recurrent saphenofemoral junction reflux after ligation and stripping of the greater saphenous vein. *J Vasc Surg* 2001;34:236-40.
  14. van Rij AM, Jones GT, Hill GB, Jiang P. Neovascularization and recurrent varicose veins: more histologic and ultrasound evidence. *J Vasc Surg* 2004;40:296-302.
  15. Stucker M, Netz K, Breuckmann F, Altmeyer P, Mumme A. Histomorphologic classification of recurrent saphenofemoral reflux. *J Vasc Surg* 2004;39:16-21.
  16. Jones L, Braithwaite BD, Selwyn D, Cooke S, Earnshaw JJ. Neovascularisation is the principal cause of varicose vein recurrence: results of a randomised trial of stripping the long saphenous vein. *Eur J Vasc Endovasc Surg* 1996;12:442-5.
  17. Stonebridge PA, Chalmers N, Beggs I, Bradbury AW, Ruckley CV. Recurrent varicose veins: a varicographic analysis leading to a new practical classification. *Br J Surg* 1995;82:60-2.
  18. Nyamekye I, Shephard NA, Davies B, Heather BP, Earnshaw JJ. Clinicopathological evidence that neovascularisation is a cause of recurrent varicose veins. *Eur J Vasc Endovasc Surg* 1998;15:412-5.
  19. Glass GM. Neovascularization in recurrence of varices of the great saphenous vein in the groin: phlebography. *Angiology* 1988;39:577-82.
  20. Chandler JG, Pichot O, Sessa C, Schuller-Petrovic S, Osse FJ, Bergan JJ. Defining the role of extended saphenofemoral junction ligation: a prospective comparative study. *J Vasc Surg* 2000;32:941-53.
  21. Merchant RF, Kistner RL. Endovenous radiofrequency obliteration of saphenous vein reflux. In: Pearce WH, Matsumura JS, Yao JS, editors. *Trends in vascular surgery 2003*. Chicago: Precept Press; 2004. p. 451-463.
  22. Merchant RF, Pichot O, Myers KA. Four-year follow-up on endovascular radiofrequency obliteration of great saphenous reflux. *Dermatol Surg* 2005;31:129-34.

Submitted Feb 11, 2005; accepted May 1, 2005.

## INVITED COMMENTARY

**Thomas W. Wakefield, MD, *Ann Arbor, Mich***

Endovenous treatment to obliterate the saphenous vein may be performed by several techniques, including radiofrequency ablation, laser treatment, or foam sclerotherapy. In this study, Dr Merchant reports the data from an ongoing multicenter prospective registry of radiofrequency ablation treatment. Clinical and duplex follow-up was performed within 1 week, at 6 months, at 1 year, and yearly thereafter. A total of 1006 patients and 1222 limbs were treated; most involved the greater saphenous vein above the knee. Vein occlusion rates ranged from 83.5% to 88.2%. Clinical symptom improvement was noted in 70% to 80% of limbs with anatomic failure and in 85% to 94% of limbs with anatomic success.

There is no question that endovenous saphenous vein ablation is here to stay. The stage was set for such a procedure by the determination that patients had better results with varicose vein surgery if the saphenous vein was removed along with saphenofemoral ligation, rather than with ligation alone.<sup>1</sup> Saphenous vein ablation, even in the presence of deep venous insufficiency, can improve the manifestations of chronic venous insufficiency.<sup>2</sup> As indicated by the authors, three randomized trials have established the superiority of endovenous radiofrequency ablation to saphenous vein surgery. The importance of the current study is to show the durability of endovenous radiofrequency closure over 5 years. Two disappointing aspects of the report include the fact that patient symptoms did not correlate with anatomic success or failure

and that body mass index correlated with anatomic failure. It is exactly in this type of patient that endovenous ablation is most appealing as opposed to operative ligation and stripping. Additionally, before radiofrequency ablation is embraced as the best method for saphenous vein ablation, more comparisons to laser and foam sclerotherapy, techniques that have shown excellent results, should be performed.<sup>3,4</sup> Nonetheless, this study is an important contribution to our understanding of the durability and efficacy of this new modality for the treatment of saphenous vein reflux.

## REFERENCES

1. Rutgers PH, Kitslaar PJ. Randomized trial of stripping versus high ligation combined with sclerotherapy in the treatment of the incompetent greater saphenous vein. *Am J Surg* 1994;168:311-5.
2. Barwell JR, Davies CE, Deacon J, Harvey K, Minor J, Sassano A, et al. Comparison of surgery and compression with compression alone in chronic venous ulceration (ESCHAR study): randomized controlled trial. *Lancet* 2004;363:1854-9.
3. Min RJ, Khilnani N, Zimmet SE. Endovenous laser treatment of saphenous vein reflux: long-term results. *J Vasc Interv Radiol* 2003;14:991-6.
4. Guex JJ. Foam sclerotherapy: an overview of use for primary venous insufficiency. *Semin Vasc Surg* 2005;18:25-29.